



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

found two of the typical three cells well developed, and each containing one large ovule. The third cell was dwarfed, and had two ovules certainly, of small size : there may have been other rudimentary ones, but not distinct enough to be made out with a good magnifying glass. I had hoped to find four cells, so as to have the tetramerism quite complete ; but in this expectation I was disappointed, partially at least.

The incident is perhaps worth recording, as the spiderwort is usually such a regular and typical trimerous flower.

D. S. MARTIN, 236, W. 4th St., July 12th.

49. *Aspidium fragrans*, Swartz,—does grow in our state. Two years ago I found a few specimens of it on the rocky precipices at Lake Avalanche in Essex County. It is not improbable, therefore, that it occurs at New Russia.

I intend visiting the Adirondack region again soon, and hope to find more of it. *Woodsia glabella*, R. Brown, another rare fern with us, also occurs there.

CHAS. H. PECK, Albany, July 6th.

50. New Stations.—*Egopodium podagraria*, L., seems to be pretty well established at Guttenberg.—*Genum strictum*, Ait., and *Erigeron Philadelphicum*, L., grow on Long Hill, N. J.—Dr. Pooley has sent us *Diospyros Virginiana*, L., from Yonkers.—We gathered in a swampy meadow at New Lots, in May, what we take to be *Rhynchospora fusca*, Rœm. & Schultes. We have little doubt of its identity though the plant was only in flower.

51. *Corrigenda*.—§ 38, read “Fern” for “Form”; § 39, read “plentifully”.

LABIATÆ.

TEUCRIUM, L.—*T. Canadense*, L.; common along shores ; N. Y.

TRICHOSTEMA, L.—*T. dichotomum*, L.; common ; N. Y.

MENTHA, L.—*M. rotundifolia*, L.; Bloomfield, N. J., and near Hudson Station, N. R. R. of N. J., *W. H. L.*—*M. viridis*, L.; common.—*M. piperita*, L.; less common, but not rare ; these two mints are not reported from N. Y. Island, but probably grow there.—*M. aquatica*, L.: Var. *crispa*, Benth.; near Greenwood Cemetery on Coney Island R. R.; Pamrapo, on Bergen Point ; and near Lake Mohegan, Westchester Co.; also in Orange Co. It is singular that this well marked variety, which seems not uncommon, should not have been noticed before. *W. H. L.*—*M. Canadensis*, L.; common ; N. Y.: Var. *glabrata*, Benth.; “wet meadows, among bushes,” *Torr. Cat.*; New Jersey, *W. H. L.*—*M. arvensis*, L., and *M. sativa*, L., are also reported, but with more or less of uncertainty, and are probably not established.

LYCOPUS, L.—*L. Virginicus*, L.: Var. *macrophyllus*; common ; N. Y.; plant intensely bitter, *Austin*; less common than the next, and not reported from Long Island, yet probably grows there.—*L. sinuatus*, Ell.; common ; N. Y.

Terms—One dollar for one copy ; five dollars for seven ; and half a dollar for every additional copy, per annum.

Local Herbarium, 3, E. 33d St.—Editor, 224, E. 10th St.